

KWAZULU NATAL TREATMENT DATA


MS JODILEE ERASMUS

Phase 57 2024b | Jul-Dec 2024





SERVICE USERS IN TREATMENT







Treatment Demand Data • Service Quality Measures (SQM) • Community-Based Harm Reduction Services

OVERVIEW OF THE STATS

Centre	2022 a	2024b
SANCA DURBAN	59	73
SANCA NEWCASTLE	58	43
SANCA NONGOMA	16	19
SANCA PIETERMARITZBURG	174	81
SANCA ZULULAND	235	138





OVERVIEW OF THE STATS CONTINUED...

Centre	2022a	2024b
MADADENI	109	0
CARELINE	22	0
NEWLANDS	39	0
ANTI-DRUG FORUM	34	0
ARCA	40	15
SERENITY ADDICTIONS	167	73





NUMBER OF SERVICE USERS IN TREATMENT







OVERVIEW OF THE STATS 18 AND UNDER



Centre	2022a	2024b
SANCA DURBAN	1	5
SANCA NEWCASTLE	10	14
SANCA NONGOMA	5	0
SANCA PIETERMARITZBURG	60	22
SANCA ZULULAND	60	30
RIVERVIEW MANOR	1	1
SERENITY ADDICTIONS	1	2
AKESO UMHLANGA	3	0
ANTI-DRUG FORUM	29	0
ARCA	1	0
MADADENI	4	0
NEWLANDS	12	0

OVERVIEW OF THE STATS





- Geographical Barriers: Are areas far from healthcare facilities?
- Difficulties with the data collection processes/ unrealistic deadline? End of Jan/End of July.
- Training needs
- Socio-Economic Factors: Affordability? Transportation and other costs associated with accessing healthcare?
- Reduced targets/ funding cuts
- Healthcare Infrastructure: limited resources, such as insufficient staff and inadequate facilities/ lack of facilities?
- Inconsistent healthcare policies in the province?






REGION	Number of cases	Number of centres
Central region	263	6
Eastern Cape	338	7
Gauteng	3199	20
Kwa Zulu Natal	483	9
Northern region	709	7
Western Cape	979	22





SACENDU SOUTH AFRICAN COMMUNITY EPIDEMIOLOGY NETWORK ON DRUG USE








SACENDU SOUTH A EDICAN COMMUNITY EDIDENIOLOGY NETWORK ON DRUG USE


Treatment Demand Data • Service Quality Measures (SQM) • Community-Based Harm Reduction Services

AGE DISTRIBUTION





HIGHEST LEVEL OF EDUCATION






SACENDU SOUTH AFRICAN COMMUNITY EPIDEMIOLOGY NETWORK ON DRUG USE

Treatment Demand Data • Service Quality Measures (SQM) • Community-Based Harm Reduction Services


EMPLOYMENT STATUS



WHERE SERVICE USERS RESIDE










WHO DO SERVICE USERS LIVE WITH












SUBSTANCES OF USE





KZN PRIMARY SUBSTANCE OF USE










SACENDU SOUTH AFRICAN COMMUNITY EPIDEMIOLOGY NETWORK ON DRUG USE






FREQUENCY OF USE



SACENDU SOUTH AFRICAN COMMUNITY EPIDEMIOLOGY NETWORK ON DRUG USE






SECONDARY SUBSTANCE OF USE




SACENDU SOUTH AFRICAN COMMUNITY EPIDEMIOLOGY NETWORK ON DRUG USE





PRIMARY SUBSTANCE OF USE BY GENDER

■ Male ■ Female ■ Other






Profile of Individuals who use first product codeine (n=44, 11%)	Percent (%)
Majority were male	84
Majority were aged: 15-19 years (32%)	
Unemployed	39%
Most common codeine product: First product: tablets, followed by syrup Second product: both syrup and tablets (n = 3)	
Type of Codeine-containing medication misused	
Stilpane; - Tablets	
Bronchleer with codeine; Benylin with codeine, Cough mixture Syrup	

SERVICE USERS REPORTING NCDS UPON ADMISSION (n = 88, 19%)





Treatment Demand Data • Service Quality Measures (SQM)




Last seen Doctor	%
Less than 2 years ago	77
2-5 years ago	17
More than 5 years ago	6

NCD BY SELECTED PRIMARY SUBSTANCE










Primary substance of use by NCD (National data: annual glance n = 2311)






Tobacco use (n = 306, 63%)





Treatment Demand Data • Service Quality Measures (SQM)








18's AND UNDER





GENDER FOR 18 YEARS AND UNDER



KZN SUMMARY



- Steady decline in numbers over the last two years
- Males dominate treatment access
- Largest age category accessing treatment is 30 34 followed by 15-19.
- Mean number of readmissions: 2
- Age category with highest proportion of readmissions: 25-29
- Most used substances: Alcohol, Dagga, Crack/Cocaine, and Heroin/Opiates
- Increase in Alcohol use, decrease in Crack/Cocaine use.
- Most common secondary substances of use: Crack/Cocaine, Alcohol, Dagga.
- Mean age for first age of primary substance use: 19
- Overall mean age for treatment access for tx: 32
- Majority codeine users: male; unemployed, largest age category reporting codeine use 15-19.
- Increase in females 18 and under accessing treatment
- Primary substances for 18 and under: Increase in Dagga and Tobacco Products; Decrease in Alcohol use





SACENDU

SOUTH AFRICAN COMMUNITY EPIDEMIOLOGY NETWORK ON DRUG USE

Treatment Demand Data • Service Quality Measures (SQM) • Community-Based Harm Reduction Services

THANK YOU BAIE DANKIE ENKOSI

For more information, contact us:

https://www.samrc.ac.za/intramural-research-units/ATOD-sacendu

jodilee.erasmus@mrc.ac.za/ nancy.hornsby@mrc.ac.za/ kamogelo.moletsane@mrc.ac.za

021 938 0313

