
# **EXAMPLE ABSTRACT**

PhD project abstract	
HIV-associated in virial adherence and management	ally suppressed people living with HIV: Challenges in
Rationale	
communicable diseases (NCDs) such as cardiovatheir HIV-negative counterparts. Increased NC inflammation, leading to endothelial dysfunction and major contributor to ongoing inflammation is detectable threshold of current viral load assays (CART) can lead to viral suppression with adherence to cause residual viraemia, significantly contributions.	currently < 20 copies/mL).  rates as low as 80%, but imperfect adherence is known ting to HIV-associated NCD development. In clinical evalent, occurring in 24%-57% of PLWH, and presents
It is currently unclear if	directly associated
with endothelial dysfunction and other HIV-associa diabetic drug with pleiotropic effects, can be repur	ted NCDs. It is also unknown if metformin, a well-knowr posed for some non-diabetic NCDs. The overall aim o les in ART adherence and address the management o
Objectives	
Adherence-as-prevention	
To determine the association of:	
1. Imperfect ART adherence and NCD development	opment in virally suppressed PLWH.
Imperfect ART adherence and	using the point-of-care EndoPAT® device.
Management of HIV-associated NCDs	
_	V-associated NCDs.
To explore the effects and mechanisms of a common NCD without effects.	as an intervention for HIV-associated sensory
<ol><li>To determine the validity and optimal dose to the drug-interaction with</li></ol>	
Prevention	Management
	Paper 5 What is the validity and optimal
Paper 1 Can	Paper 3 What is the validity and optimal
Is imperfect ART adherence	
pre	went or treat HIV-
1	
Imperfect adherence Chronic inflammation & endothelial dysfunction	HIV-associated non- Drug interactions communicable diseases
Figure 1: Doctoral themes,	
coherence, and Paper 2	♥ Paper 4
research questions Is endothelial dysfunction using	
the p	changes in epidermal nerve fibre

dev....unsuppressed HIV

a function of ART adherence?



Retrospective, longitudinal cohort study, using existing data from the longitudinal AIDS Clinical Trials Group (ACTG) research cohorts I

# Paper 2:

Case-control sub-study of the active care ART adherence device.

testing a

# Paper 3:

Scoping review using a predefined search strategy in multiple electronic databases.

# Paper 4:

Exploratory, pilot cohort sub-study of the active \_\_\_\_\_\_ that is assessing PLWH with metabolic syndrome.

# Paper 5:

Pharmacokinetic drug-interaction study using directly-observed metformin and dolutegravir administration with intensive metformin pharmacokinetic blood sampling and genetic drug transporter analysis.

#### Feasibility

All the studies of the PhD project have high feasibility of successful completion within the allocated time. The feasibility of all the PhD studies is ensured by the study designs and secured funding:

- Paper 1 is conducted as a longitudinal study of existing data after study completion has already occurred, and Paper 2 as a retrospective case-control design at the last visit of the parent study, with low anticipated drop-out. Additionally, Paper 2 will only seek to enrol approximately 80% of the parent study population, significantly improving the likelihood of successful completion.
- Paper 2, 4, and 5 are linked to existing parent studies that are active and recruiting, with the PIs having strong track records of successful previous study completions.
- Paper 1, 2, 4, and 5 have good probability of reaching their respective sample sizes, due to the experienced teams conducting the parent studies and successful environments of these studies.
- At the time of this application, the prospective data collection studies (Papers 2, 4, and 5) have already reached between 20% 50% of their sample sizes.
- Funding for all the PhD studies has been secured for all objectives and for the duration of the studies and overall PhD project.

#### Research translation & dissemination plan

The research will be translated into non-scientific language and shared with the communities from where it was obtained, to include participants and communities throughout the research process. Research findings will be disseminated through presentations at local and international scientific meetings, accredited journal publications, institutional media releases, and by placing raw data in a protected data repository for potential future use.

#### **Potential impact**

The research findings will add valuable data on NCD associations in virally suppressed PLWH, the potential of the versatile drug metformin to be used in HIV care, and

PLWH. This has significant immediate and future implications for HIV-associated NCDs in South Africa and globally.