## REPORT ON WEEKLY DEATHS IN SOUTH AFRICA

# 9 – 15 OCTOBER 2022 (WEEK 41)

Debbie Bradshaw, Ria Laubscher,
Rob Dorrington, Pam Groenewald, Tom Moultrie

Burden of Disease Research Unit South African Medical Research Council 18 October 2022





#### Glossary:

Age-standardised excess death rate: Indirectly age-standardised excess death rates have been calculated for each province to adjust the crude death rates per capita for the differences in distribution of the population by age. The adjustment factor for each province is calculated as the crude death rate for South Africa divided by what the crude rate for South Africa would have been had the age distribution of the population been that of the province. Standardisation for age is necessary when comparing populations that differ in their age structure because age has a powerful influence on the risk of dying. The rate is based on the cumulative number of excess deaths since 3 May 2020 to date divided by the population estimate for 2021 and has not been annualised.

Actual number of deaths: The actual number of deaths in South Africa have been estimated from the numbers recorded on the National Population Register using weighting factors set to produce results consistent with those of the annual Rapid Mortality Surveillance Report to account for deaths of persons who are not on the National Population Register as well as those that have not been registered with the Department of Home Affairs. The adjustments to account for incompleteness of recording of deaths on the NPR have been re-estimated for the 2021 reports taking into account the 2017 cause-of-death data released by Stats SA in 2020. A methodological note briefly outlining the changes can be downloaded with this report from the SAMRC website: <a href="https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa">https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa</a>.

**Epi-week:** The Weekly Death Reports in 2020 used weeks from 1 January and ran from Wednesday to Tuesday. In setting up the monitoring for 2021, we recast the data to report by an 'Epi-week' consistent with CDC and many NICD reports which run from Sunday to Saturday, ensuring continuity of weeks from one year to the next. Each week is aligned with the 'Epi-year' that has 4 or more days in that week. Week 53 of 2020 is from 27 December 2020 to 2 January 2021, Week 1 of 2021 is 3 January – 9 January 2021 and Week 1 of 2022 is 2 January – 8 January 2022.

**Excess deaths:** There is no universal definition of, or understanding of what is meant by, "excess mortality". It is a term used in epidemiology and public health that refers to the number of deaths that are occurring above what we would normally expect. The WHO uses the term to describe "Mortality above what would be expected based on the non-crisis mortality rate in the population of interest. Excess mortality is thus mortality that is attributable to the crisis conditions. It can be expressed as a rate (the difference between observed and non-crisis mortality rates), or as a total number of excess deaths."

Excess natural deaths associated with COVID-19: Generally, the number of excess deaths per week is calculated as the number of all-cause deaths in that week less the number that might be assumed to have occurred had there not been the epidemic (i.e. the counterfactual number), provided that the counterfactual is lower. However, this approach has generally only been applied to countries where deaths have been tracking the counterfactual before the onset of significant numbers of COVID-19 related deaths. The method provides a poor estimate of the numbers of COVID-19 and collateral deaths in the early stages of the epidemic when this is not the case. Thus, we estimated the numbers of COVID-19 and collateral deaths, once a clear upward trend is evident, as the number of actual deaths less a baseline number determined as a proportion of the predicted number. By the end of the 1st wave of the pandemic, the predicted values have been used as the counterfactual.

**General warning:** The Department of Home Affairs has faced sporadic temporary office closures, particularly in areas that are more affected by COVID-19. This may affect our allocation of deaths to a particular metro area. For example, a death that occurred in the City of Cape Town might have been registered at an office outside of the City because of a temporary closure of offices in the metro. Closure may also cause a delay in the processing of the death registration which would result in an underestimate of the deaths in the most recent week.

### **Background**

This report provides estimates of the weekly number of deaths of all persons in South Africa for epidemiological **Week 41** of 2022, covering the period **9 – 15 October 2022. Warning:** Estimates for the most recent few weeks need to be treated with caution.

The predicted numbers of weekly deaths in 2022 have been estimated using the models prepared for 2021 with population estimates for 2022. A methodological note outlining the approach for monitoring deaths during 2021 can be downloaded from the SAMRC website: <a href="https://www.samrc.ac.za/sites/default/files/files/2021-01-24/Methodological Note on Predicted Weekly Deaths 20 Jan 2021.pdf">https://www.samrc.ac.za/sites/default/files/files/2021-01-24/Methodological Note on Predicted Weekly Deaths 20 Jan 2021.pdf</a>.

Briefly, the predicted values for the provinces and nationally are based on negative binomial models based on death data for the period 2014-2019. After reviewing trends in the data, separate negative binomial models have been fitted to the unnatural deaths for all provinces combined, and separate models for natural deaths for each of KwaZulu-Natal, Western Cape and the Northern Cape, and the 6 other provinces in a combined model to provide estimates by age, sex and epi-week for each year. A prediction interval has been estimated on the basis of the variability in the observed weekly data for each reported domain. Predicted values for each metropolitan area continue to be based on the time-series model of data for 2018 and 2019 developed for 2020 as the trends in the sub-provincial data need further investigation to develop a comprehensive district-level model.

#### **Trends**


- The weekly number of deaths (all ages) from all causes has been tracking just above the upper prediction bound for several weeks and increased to 10,832 deaths in Week 41 (9 15 Oct 2022), associated with an increase in deaths from both natural and unnatural causes. Warning: Estimates for the most recent few weeks need to be treated with caution.
- The number of deaths from natural causes (all ages) has been running just below the upper prediction bound and increased to 1,161 excess natural deaths in Week 41 (9 15 Oct 2022).
- Since 3 May 2020, there has been a cumulative total of about **334,000** excess deaths from natural causes of persons all ages of which **85,000** occurred in 2020, **203,200** occurred in 2021 (since 3 Jan 2021) and **47,400** have occurred in 2022 by the end of **Week 41** (9 15 Oct 2022).

Week	Date	Weekly excess deaths from natural causes (all ages)	Cumulative excess since 3 May 2020 (all ages)	Cumulative excess since 2 January 2022 (all ages)
31	31-Jul-22 – 6-Aug-22	772	326,500	39,570
32	7-Aug-22 – 13-Aug-22	501	327,001	40,071
33	14-Aug-22 – 20-Aug-22	723	327,724	40,794
34	21-Aug-22 – 27-Aug-22	846	328,570	41,640
35	28-Aug-22 – 3-Sep-22	841	329,412	42,482
36	4-Sep-22 – 10-Sep-22	861	330,273	43,343
37	11-Sep-22 – 17-Sep-22	634	330,908	43,978
38	18-Sep-22 – 24-Sep-22	685	331,593	44,663
39	25-Sep-22 – 1-Oct-22	763	332,356	45,426
40	2-Oct-22 – 8-Oct-22	863	333,220	46,290
41	9-Oct-22 – 15-Oct-22	1,163	334,383	47,453

- For people under the age of 60, the number of natural deaths has been tracking close to the predicted value since mid-June. The cumulative number of excess natural deaths for people under-60 years since **3 May 2020** was about **86,600**.
- The number of weekly excess deaths in the 60+ years age group has continued to track just above the upper prediction bound. In **Week 41** (9 15 Oct 2022), there was an increase to **845** excess deaths with a p-score of 21%. The cumulative total number of excess natural deaths in this age group since **3 May 2020** is about **247,700**.

Week	Date	Weekly excess deaths from natural causes for persons 60+ years	p-score
31	31-Jul-22 – 6-Aug-22	773	17.3%
32	7-Aug-22 – 13-Aug-22	538	12.0%
33	14-Aug-22 – 20-Aug-22	637	14.4%
34	21-Aug-22 – 27-Aug-22	732	17.0%
35	28-Aug-22 – 3-Sep-22	639	15.1%
36	4-Sep-22 – 10-Sep-22	596	13.7%
37	11-Sep-22 – 17-Sep-22	577	13.7%
38	18-Sep-22 – 24-Sep-22	583	14.2%
39	25-Sep-22 – 1-Oct-22	753	18.7%
40	2-Oct-22 – 8-Oct-22	752	17.9%
41	9-Oct-22 – 15-Oct-22	845	20.9%

- Although caution is required not to overinterpret the estimates for the most recent weeks, the natural deaths experienced an uptick in **Week 40** (2 8 Oct 2022) in several provinces. And while the numbers in **Gauteng** and **Western Cape** may have reversed in **Week 41** (9 15 Oct 2022), this is not the case in **Eastern Cape**, **Limpopo** and **KwaZulu Natal** where they have continued to increase slightly.
- Per capita excess death rates have been calculated for the provinces to scale the cumulative deaths for the population size of each province (**Table 1**). By the end of **Week 41** (9 15 Oct 2022), the national excess death rate since 3 May 2020 was **562** per **100,000** population.
- The provinces with the highest cumulative numbers of excess deaths at the end of Week 41 (9 15 Oct 2022), are, in order, KwaZulu-Natal, Gauteng and Eastern Cape. The ranking changes to Eastern Cape, Northern Cape and Free State for the crude death rates per capita (i.e., taking size of the provincial populations into account) and to Northern Cape, Eastern Cape and KwaZulu-Natal using the agestandardised rates per capita (i.e., taking into account the age distribution of the provincial population). The Western Cape, followed by Gauteng, continue to have the lowest cumulative age standardised per capita rates.
- The number of unnatural deaths reached a high of **1,557** in **Week 15** (10 16 Apr 2022), the week of the floods in KwaZulu-Natal, and reverted to the predicted monthly cyclical pattern thereafter. After tracking within the prediction bounds for several weeks, the number of unnatural deaths in **Week 41** (9 15 Oct 2022) has shown an uptick with **1,450** deaths.



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations


Table 1: Number of excess natural deaths of persons by province and metro relative to revised predicted number based on the observed drop during lockdown, South Africa 2020/22


Region	Period	Excess deaths vs revised base	Excess deaths per 100,000 population	Age standardised excess death rate per 100,000
South Africa	3 May 20 – 15 Oct 22	334,383	562	562
Province				
Eastern Cape	31 May 20 – 15 Oct 22	57,974	881	710
Free State	21 Jun 20 – 15 Oct 22	19,315	663	663
Gauteng	7 Jun 20 – 15 Oct 22	65,918	423	465
KwaZulu-Natal	7 Jun 20 – 15 Oct 22	68,671	600	690
Limpopo	21 Jun 20 – 15 Oct 22	37,189	629	551
Mpumalanga	21 Jun 20 – 15 Oct 22	25,771	536	577
Northern Cape	28 Jun 20 – 15 Oct 22	10,169	869	813
North West	28 Jun 20 – 15 Oct 22	18,267	454	466
Western Cape	3 May 20 – 15 Oct 22	31,107	441	388
Metropolitan Municipa	ality			
Buffalo City	31 May 20 – 15 Oct 22	7,407		
City of Cape Town	3 May 20 – 15 Oct 22	24,209		
Ekurhuleni	7 Jun 20 – 15 Oct 22	14,909		
eThekwini	14 Jun 20 – 15 Oct 22	14,086		
Johannesburg	7 Jun 20 – 15 Oct 22	22,820		
Mangaung	21 Jun 20 – 15 Oct 22	6,045		
Nelson Mandela Bay	31 May 20 – 15 Oct 22	8,908		
City of Tshwane	7 Jun 20 – 15 Oct 22	12,075		


Note: Period has been determined based on when an upturn in the number of natural deaths became apparent. Parts do not sum to the whole because office closures due to Covid-19 may have led to registration of deaths at other offices which may not be in the same area, and random fluctuation at the point at which the baseline is determined.

Table 2: Number of excess deaths from all causes of persons by province and metro relative to predicted number based on historical trend, South Africa 2020/22


2	Excess deaths vs	Excess deaths per	
Region	forecast	100,000 population	
South Africa	339,172	570	
Province			
Eastern Cape	60,934	926	
Free State	19,168	658	
Gauteng	63,195	405	
KwaZulu-Natal	72,556	634	
Limpopo	37,808	640	
Mpumalanga	25,656	533	
Northern Cape	10,117	864	
North West	17,720	440	
Western Cape	32,017	454	
Metropolitan Municipality			
Buffalo City	8,018		
City of Cape Town	22,937		
Ekurhuleni	13,729		
eThekwini	13,216		
Johannesburg	23,332		
Mangaung	6,900		
Nelson Mandela Bay	9,054		
City of Tshwane	12,698		











Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations


















































Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations. As only a quarter to a third of unnatural deaths in the most recent week are processed at the time of the survey, the estimate for the most recent week is quite uncertain.

#### Vertical lines in order

- 0 Week Disaster Management Act implemented
- 1 Week lockdown level 5 introduced
- 2 Week lockdown changed to level 4, with curfew
- 3 Week lockdown changed to level 3 including unbanning of alcohol
- 4 Week alcohol re-banned, and a curfew re-introduced
- 5 Week lockdown changed to level 2, including unbanning of alcohol
- 6 Week lockdown changed to level 1
- Week lockdown changed to level 3 advanced (re-banning alcohol and a extension of curfew)
- 8 Week lockdown relaxed to allow sale of alcohol 4 days/week and reduce curfew
- 9 Week lockdown relaxed to allow sale of alcohol except during curfew and reduce curfew to midnight to 4am
- 10 Week lockdown changed to level 3 advanced (limiting alcohol and a extending of curfew)
- 11 Week lockdown changed to level 4, with re-banning of alcohol, curfew 9pm-4am
- 12 Week of unrest in KZN and GT
- 13 Week lockdown changed to level 3 advanced (alcohol 4 days/w, curfew 10pm-4am)
- 14 Week lockdown changed to level 2 advanced (alcohol 5 days/w, curfew 11pm-4am)
- 15 Week lockdown changed to level 1 advanced (no alcohol post 11pm, curfew 12pm-4am, large gatherings)
- 16 Week lockdown level 1 advanced (removed limits on alcohol & curfew, allowed larger gatherings)
- 17 Week lockdown level 1 advanced (allowed full school attendance, reduced isolation & quarantine requirements)
- 18 Week lockdown level 1 advanced (no masks outdoors, larger gathering permitted, no travel testing for vaccinated)
- 19 KZN floods
- 20 All COVID restrictions lifted on 22 June