## REPORT ON WEEKLY DEATHS IN SOUTH AFRICA

# 10 – 16 JULY 2022 (WEEK 28)

Debbie Bradshaw, Ria Laubscher, Rob Dorrington, Pam Groenewald, Tom Moultrie

Burden of Disease Research Unit South African Medical Research Council 19 July 2022



#### Glossary:

**Age-standardised excess death rate:** Indirectly age-standardised excess death rates have been calculated for each province to adjust the crude death rates per capita for the differences in distribution of the population by age. The adjustment factor for each province is calculated as the crude death rate for South Africa divided by what the crude rate for South Africa would have been had the age distribution of the population been that of the province. Standardisation for age is necessary when comparing populations that differ in their age structure because age has a powerful influence on the risk of dying. The rate is based on the cumulative number of excess deaths since 3 May 2020 to date divided by the population estimate for 2021 and has not been annualised.

Actual number of deaths: The actual number of deaths in South Africa have been estimated from the numbers recorded on the National Population Register using weighting factors set to produce results consistent with those of the annual Rapid Mortality Surveillance Report to account for deaths of persons who are not on the National Population Register as well as those that have not been registered with the Department of Home Affairs. The adjustments to account for incompleteness of recording of deaths on the NPR have been re-estimated for the 2021 reports taking into account the 2017 cause-of-death data released by Stats SA in 2020. A methodological note briefly outlining the changes can be downloaded with this report from the SAMRC website: <a href="https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa">https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa</a>.

**Epi-week:** The Weekly Death Reports in 2020 used weeks from 1 January and ran from Wednesday to Tuesday. In setting up the monitoring for 2021, we recast the data to report by an 'Epi-week' consistent with CDC and many NICD reports which run from Sunday to Saturday, ensuring continuity of weeks from one year to the next. Each week is aligned with the 'Epi-year' that has 4 or more days in that week. Week 53 of 2020 is from 27 December 2020 to 2 January 2021, Week 1 of 2021 is 3 January – 9 January 2021 and Week 1 of 2022 is 2 January – 8 January 2022.

**Excess deaths:** There is no universal definition of, or understanding of what is meant by, "excess mortality". It is a term used in epidemiology and public health that refers to the number of deaths that are occurring above what we would normally expect. The WHO uses the term to describe "Mortality above what would be expected based on the non-crisis mortality rate in the population of interest. Excess mortality is thus mortality that is attributable to the crisis conditions. It can be expressed as a rate (the difference between observed and non-crisis mortality rates), or as a total number of excess deaths."

**Excess natural deaths associated with COVID-19**: Generally, the number of excess deaths per week is calculated as the number of all-cause deaths in that week less the number that might be assumed to have occurred had there not been the epidemic (i.e. the counterfactual number), provided that the counterfactual is lower. However, this approach has generally only been applied to countries where deaths have been tracking the counterfactual before the onset of significant numbers of COVID-19 related deaths. The method provides a poor estimate of the numbers of COVID-19 and collateral deaths in the early stages of the epidemic when this is not the case. Thus, we estimated the numbers of COVID-19 and collateral deaths, once a clear upward trend is evident, as the number of actual deaths less a baseline number determined as a proportion of the predicted number. By the end of the 1<sup>st</sup> wave of the pandemic, the predicted values have been used as the counterfactual.

**General warning:** The Department of Home Affairs has faced sporadic temporary office closures, particularly in areas that are more affected by COVID-19. This may affect our allocation of deaths to a particular metro area. For example, a death that occurred in the City of Cape Town might have been registered at an office outside of the City because of a temporary closure of offices in the metro. Closure may also cause a delay in the processing of the death registration which would result in an underestimate of the deaths in the most recent week.

### Background

This report provides estimates of the weekly number of deaths of all persons in South Africa for epidemiological **Week 28** of 2022, covering the period **10 – 16 July 2022.** 

**Warning:** There is added uncertainty about the estimates of the number of deaths in most recent week. Data received from the Department of Home Affairs on 19 July 2022 indicated that no deaths had been processed on the National Population Register on the previous day. An adjustment has been made to estimate the number of deaths that have occurred but not yet processed, treating the data as if the 18 July had been a public holiday, however, it is not clear when the hiatus in processing began, so the number of natural deaths in the most recent week may be underestimated.

The predicted numbers of weekly deaths in 2022 have been estimated using the models prepared for 2021 with population estimates for 2022. A methodological note outlining the approach for monitoring deaths during 2021 can be downloaded from the SAMRC website: <u>https://www.samrc.ac.za/sites/default/files/files/2021-01-</u>24/Methodological Note on Predicted Weekly Deaths 20 Jan 2021.pdf.

Briefly, the predicted values for the provinces and nationally are based on negative binomial models based on death data for the period 2014-2019. After reviewing trends in the data, separate negative binomial models have been fitted to the unnatural deaths for all provinces combined, and separate models for natural deaths for each of KwaZulu-Natal, Western Cape and the Northern Cape, and the 6 other provinces in a combined model to provide estimates by age, sex and epi-week for each year. A prediction interval has been estimated on the basis of the variability in the observed weekly data for each reported domain. Predicted values for each metropolitan area continue to be based on the timeseries model of data for 2018 and 2019 developed for 2020 as the trends in the sub-provincial data need further investigation to develop a comprehensive district-level model.

### Trends

- The weekly number of deaths (all ages) from all causes was 11,011 deaths in Week 28 (10 16 Jul 2022) and falls within the prediction bounds. Warning: There is added uncertainty about the number of deaths in Week 28 as no deaths were processed on the National Population Register on Monday 18 July.
- The number of excess deaths from natural causes (all ages) increased to a high of 2,032 in Week 21 (22 28 May 2022) and appear to have decreased to 448 in Week 28 (10 16 Jul 2022). Warning: There is added uncertainty about the number of deaths in Week 28 as no deaths were processed on the National Population Register on Monday 18 July.
- Since 3 May 2020, there has been a cumulative total of about 325,000 excess deaths from natural causes of persons all ages of which 85,000 occurred in 2020, 203,200 occurred in 2021 (since 3 Jan 2021) and 37,000 have occurred in 2022 by the end of Week 28 (10 16 Jul 2022).

Week	Date	Weekly excess deaths from natural causes (all ages)	Cumulative excess since 3 May 2020 (all ages)	Cumulative excess since 2 January 2022 (all ages)
21	22-May-22 – 28-May-22	2,032	317,264	28,865
22	29-May-22 – 4-Jun-22	1,668	318,932	30,533
23	5-Jun-22 – 11-Jun-22	1,353	320,284	31,886
24	12-Jun-22 – 18-Jun-22	1,431	321,715	33,316
25	19-Jun-22 – 25-Jun-22	1,152	322,867	34,468

26	26-Jun-22 – 2-Jul-22	1,191	324,058	35,659
27	3-Jul-22 – 9-Jul-22	844	324,902	36,503
28	10-Jul-22 – 16-Jul-22	448	325,350	36,951

- For people under the age of 60, the number of natural deaths was tracking between the predicted value and the upper prediction bound with a clear increase to the upper prediction bound in Week 18 (1 7 May 2022) and Week 19 (8 14 May 2022) and decreased slightly thereafter. The cumulative number of excess natural deaths for people under-60 years since 3 May 2020 was about 85,700.
- The number of weekly excess deaths in the 60+ years age group had decreased to 423 in Week 11 (13 19 Mar 2022) with a p-score of 11% and increased to 1,499 in Week 21 (22 28 May 2022) with a p-score of 36% and then decreased to 717 in Week 28 (10 16 Jul 2022) with a p-score of 15%. The cumulative total number of excess natural deaths in this age group since 3 May 2020 is just over 239,700.

Week	Date	Weekly excess deaths from natural causes for persons 60+ years	p-score
21	22-May-22 – 28-May-22	1,499	36.2%
22	29-May-22 – 4-Jun-22	1,248	28.2%
23	5-Jun-22 – 11-Jun-22	1,077	22.8%
24	12-Jun-22 – 18-Jun-22	1,275	26.6%
25	19-Jun-22 – 25-Jun-22	1,126	23.8%
26	26-Jun-22 – 2-Jul-22	1,092	23.2%
27	3-Jul-22 – 9-Jul-22	919	19.4%
28	10-Jul-22 – 16-Jul-22	717	15.4%

- KwaZulu-Natal and Limpopo have joined Mpumalanga, North West and Western Cape provinces to track, close to their predicted number in Week 28 (10 16 Jul 2022), while number of natural deaths in the provinces continued tracking along the upper prediction bound.
- Per capita excess death rates have been calculated for the provinces to scale the cumulative deaths for the population size of each province (**Table 1**). By the end of **Week 28** (10 16 Jul 2022), the national excess death rate since 3 May 2020 was **547** per **100,000** population.
- The provinces with the highest cumulative numbers of excess deaths at the end of Week 27 (3 9 Jul 2022), are, in order, KwaZulu-Natal, Gauteng and Eastern Cape. The ranking changes to Eastern Cape, Northern Cape and Free State for the crude death rates per capita (i.e., taking size of the provincial populations into account) and to Northern Cape, Eastern Cape and KwaZulu-Natal using the age-standardised rates per capita (i.e., taking into account the age distribution of the provincial population). The Western Cape, followed by Gauteng, continue to have the lowest cumulative age standardised per capita rates.
- The number of unnatural deaths reached a high of 1,547 in Week 15 (10 16 Apr 2022), the week of the floods in KwaZulu-Natal, and has resumed the monthly cyclical pattern with 1,305 deaths in Week 28 (10 16 Jul 2022), tending to oscillate between the predicted number during the trough and the upper prediction bound at the peak. Warning: There is added uncertainty about the number of deaths in Week 28 as no deaths were processed on the National Population Register on Monday 18 July.





Table 1: Number of excess natural deaths of persons by province and metro relative to revised predicted number based on the observed drop during lockdown, South Africa 2020/22

Region	Period	Excess deaths vs revised base	Excess deaths per 100,000 population	Age standardised excess death rate per 100,000
South Africa	3 May 20 – 16 Jul 22	325,350	547	547
Province				
Eastern Cape	31 May 20 – 16 Jul 22	55,662	846	682
Free State	21 Jun 20 – 16 Jul 22	18,650	641	641
Gauteng	7 Jun 20 – 16 Jul 22	64,340	412	454
KwaZulu-Natal	7 Jun 20 – 16 Jul 22	66,435	580	668
Limpopo	21 Jun 20 – 16 Jul 22	35,574	602	527
Mpumalanga	21 Jun 20 – 16 Jul 22	24,895	517	557
Northern Cape	28 Jun 20 – 16 Jul 22	9,770	835	781
North West	28 Jun 20 – 16 Jul 22	17,972	446	458
Western Cape	3 May 20 – 16 Jul 22	32,052	454	400
Metropolitan Municipality				
Buffalo City	31 May 20 – 16 Jul 22	6,677		
City of Cape Town	3 May 20 – 16 Jul 22	23,211		
Ekurhuleni	7 Jun 20 – 16 Jul 22	14,802		
eThekwini	14 Jun 20 – 16 Jul 22	13,878		
Johannesburg	7 Jun 20 – 16 Jul 22	22,089		
Mangaung	21 Jun 20 – 16 Jul 22	5,534		
Nelson Mandela Bay	31 May 20 – 16 Jul 22	8,529		
City of Tshwane	7 Jun 20 – 16 Jul 22	11,888		


Note: Period has been determined based on when an upturn in the number of natural deaths became apparent. Parts do not sum to the whole because office closures due to Covid-19 may have led to registration of deaths at other offices which may not be in the same area, and random fluctuation at the point at which the baseline is determined.


Table 2: Number of excess deaths from all causes of persons by province and metro relative to predictednumber based on historical trend, South Africa 2020/22

Decien	Excess deaths vs	Excess deaths per	
Region	forecast	100,000 population	
South Africa	325,995	548	
Province			
Eastern Cape	57,854	879	
Free State	18,468	634	
Gauteng	61,566	395	
KwaZulu-Natal	69,309	605	
Limpopo	36,006	609	
Mpumalanga	24,788	515	
Northern Cape	9,686	828	
North West	17,436	433	
Western Cape	30,882	437	
Metropolitan Municipality			
Buffalo City	7,116		
City of Cape Town	21,815		
Ekurhuleni	13,631		
eThekwini	12,991		
Johannesburg	22,385		
Mangaung	6,254		
Nelson Mandela Bay	8,617		
City of Tshwane	12,349		





Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations








Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations




















Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations. As only a quarter to a third of unnatural deaths in the most recent week are processed at the time of the survey, the estimate for the most recent week is quite uncertain.

#### Vertical lines in order

- 0 Week Disaster Management Act implemented
- 1 Week lockdown level 5 introduced
- 2 Week lockdown changed to level 4, with curfew
- 3 Week lockdown changed to level 3 including unbanning of alcohol
- 4 Week alcohol re-banned, and a curfew re-introduced
- 5 Week lockdown changed to level 2, including unbanning of alcohol
- 6 Week lockdown changed to level 1
- 7 Week lockdown changed to level 3 advanced (re-banning alcohol and a extension of curfew)
- 8 Week lockdown relaxed to allow sale of alcohol 4 days/week and reduce curfew
- 9 Week lockdown relaxed to allow sale of alcohol except during curfew and reduce curfew to midnight to 4am
- 10 Week lockdown changed to level 3 advanced (limiting alcohol and a extending of curfew)
- 11 Week lockdown changed to level 4, with re-banning of alcohol, curfew 9pm-4am
- 12 Week of unrest in KZN and GT
- 13 Week lockdown changed to level 3 advanced (alcohol 4 days/w, curfew 10pm-4am)
- 14 Week lockdown changed to level 2 advanced (alcohol 5 days/w, curfew 11pm-4am)
- 15 Week lockdown changed to level 1 advanced (no alcohol post 11pm, curfew 12pm-4am, large gatherings)
- 16 Week lockdown level 1 advanced (removed limits on alcohol & curfew, allowed larger gatherings)
- 17 Week lockdown level 1 advanced (allowed full school attendance, reduced isolation & quarantine requirements)
- 18 Week lockdown level 1 advanced (no masks outdoors, larger gathering permitted, no travel testing for vaccinated)
- 19 KZN floods