# REPORT ON WEEKLY DEATHS IN SOUTH AFRICA

# 20 – 26 NOVEMBER 2022 (WEEK 47)

Debbie Bradshaw, Ria Laubscher,
Rob Dorrington, Pam Groenewald, Tom Moultrie

Burden of Disease Research Unit South African Medical Research Council 29 November 2022



## Glossary:

Age-standardised excess death rate: Indirectly age-standardised excess death rates have been calculated for each province to adjust the crude death rates per capita for the differences in distribution of the population by age. The adjustment factor for each province is calculated as the crude death rate for South Africa divided by what the crude rate for South Africa would have been had the age distribution of the population been that of the province. Standardisation for age is necessary when comparing populations that differ in their age structure because age has a powerful influence on the risk of dying. The rate is based on the cumulative number of excess deaths since 3 May 2020 to date divided by the population estimate for 2021 and has not been annualised.

Actual number of deaths: The actual number of deaths in South Africa have been estimated from the numbers recorded on the National Population Register using weighting factors set to produce results consistent with those of the annual Rapid Mortality Surveillance Report to account for deaths of persons who are not on the National Population Register as well as those that have not been registered with the Department of Home Affairs. The adjustments to account for incompleteness of recording of deaths on the NPR have been re-estimated for the 2021 reports taking into account the 2017 cause-of-death data released by Stats SA in 2020. A methodological note briefly outlining the changes can be downloaded with this report from the SAMRC website: <a href="https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa">https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa</a>.

**Epi-week:** The Weekly Death Reports in 2020 used weeks from 1 January and ran from Wednesday to Tuesday. In setting up the monitoring for 2021, we recast the data to report by an 'Epi-week' consistent with CDC and many NICD reports which run from Sunday to Saturday, ensuring continuity of weeks from one year to the next. Each week is aligned with the 'Epi-year' that has 4 or more days in that week. Week 53 of 2020 is from 27 December 2020 to 2 January 2021, Week 1 of 2021 is 3 January – 9 January 2021 and Week 1 of 2022 is 2 January – 8 January 2022.

**Excess deaths:** There is no universal definition of, or understanding of what is meant by, "excess mortality". It is a term used in epidemiology and public health that refers to the number of deaths that are occurring above what we would normally expect. The WHO uses the term to describe "Mortality above what would be expected based on the non-crisis mortality rate in the population of interest. Excess mortality is thus mortality that is attributable to the crisis conditions. It can be expressed as a rate (the difference between observed and non-crisis mortality rates), or as a total number of excess deaths."

Excess natural deaths associated with COVID-19: Generally, the number of excess deaths per week is calculated as the number of all-cause deaths in that week less the number that might be assumed to have occurred had there not been the epidemic (i.e. the counterfactual number), provided that the counterfactual is lower. However, this approach has generally only been applied to countries where deaths have been tracking the counterfactual before the onset of significant numbers of COVID-19 related deaths. The method provides a poor estimate of the numbers of COVID-19 and collateral deaths in the early stages of the epidemic when this is not the case. Thus, we estimated the numbers of COVID-19 and collateral deaths, once a clear upward trend is evident, as the number of actual deaths less a baseline number determined as a proportion of the predicted number. By the end of the 1<sup>st</sup> wave of the pandemic, the predicted values have been used as the counterfactual.

**General warning:** The Department of Home Affairs has faced sporadic temporary office closures, particularly in areas that are more affected by COVID-19. This may affect our allocation of deaths to a particular metro area. For example, a death that occurred in the City of Cape Town might have been registered at an office outside of the City because of a temporary closure of offices in the metro. Closure may also cause a delay in the processing of the death registration which would result in an underestimate of the deaths in the most recent week.

# **Background**

This report provides estimates of the weekly number of deaths of all persons in South Africa for epidemiological **Week 47** of 2022, covering the period **20 – 26 November 2022. Warning:** Estimates for the most recent few weeks need to be treated with caution.

The predicted numbers of weekly deaths in 2022 have been estimated using the models prepared for 2021 with population estimates for 2022. A methodological note outlining the approach for monitoring deaths during 2021 can be downloaded from the SAMRC website: <a href="https://www.samrc.ac.za/sites/default/files/files/2021-01-24/Methodological Note on Predicted Weekly Deaths 20 Jan 2021.pdf">https://www.samrc.ac.za/sites/default/files/files/2021-01-24/Methodological Note on Predicted Weekly Deaths 20 Jan 2021.pdf</a>.

Briefly, the predicted values for the provinces and nationally are based on negative binomial models based on death data for the period 2014-2019. After reviewing trends in the data, separate negative binomial models have been fitted to the unnatural deaths for all provinces combined, and separate models for natural deaths for each of KwaZulu-Natal, Western Cape and the Northern Cape, and the 6 other provinces in a combined model to provide estimates by age, sex and epi-week for each year. A prediction interval has been estimated on the basis of the variability in the observed weekly data for each reported domain. Predicted values for each metropolitan area continue to be based on the time-series model of data for 2018 and 2019 developed for 2020 as the trends in the sub-provincial data need further investigation to develop a comprehensive district-level model.

### **Trends**

- Notification: The final weekly death report for 2022 will be released on Wed 13 December. From next year, we will no longer be reporting on excess deaths on a weekly basis. However, we plan to monitor the weekly deaths and, provided there are no unusual trends, plan to release a monthly report on the weekly numbers of deaths.
- After tracking just above the upper prediction bound for several weeks, the weekly number of deaths (all ages) from all causes tracked within the prediction bound from Week 43 (23 29 Oct 2022) till Week 46 (13 19 Nov 2022). The number was 9,434 deaths in Week 47 (20 26 Nov 2022), just above the upper prediction bound. Warning: Estimates for the most recent few weeks need to be treated with caution.
- The number of excess deaths from natural causes (all ages) increased to **982** in **Week 40** (2 8 Oct 2022) and has dropped to **536** in **Week 47** (20 26 Nov 2022).
- Since 3 May 2020, there has been a cumulative total of about **337,700** excess deaths from natural causes of persons all ages of which **85,000** occurred in 2020, **203,200** occurred in 2021 (since 3 Jan 2021) and **50,775** have occurred in 2022 by the end of **Week 47** (20 26 Nov 2022).


Week	Date	Weekly excess deaths from	Cumulative excess since	Cumulative excess since
		natural causes	3 May 2020	2 January 2022
		(all ages)	(all ages)	(all ages)
37	11-Sep-22 – 17-Sep-22	654	331,153	44,181
38	18-Sep-22 – 24-Sep-22	713	331,866	44,894
39	25-Sep-22 – 1-Oct-22	820	332,685	45,713
40	2-Oct-22 – 8-Oct-22	982	333,667	46,695
41	9-Oct-22 – 15-Oct-22	842	334,510	47,538
42	16-Oct-22 – 22-Oct-22	475	334,984	48,012
43	23-Oct-22 – 29-Oct-22	330	335,315	48,343

44	30-Oct-22 – 5-Nov-22	618	335,932	48,960
45	6-Nov-22 – 12-Nov-22	567	336,500	49,528
46	13-Nov-22 – 19-Nov-22	712	337,211	50,239
47	20-Nov-22 – 26-Nov-22	536	337,747	50,775

- For people under the age of 60, the number of natural deaths has been tracking close to the predicted value since mid-June 2022. The cumulative number of excess natural deaths for people under-60 years since **3 May 2020** was about **86,600**.
- The number of weekly excess deaths in the 60+ years age group has continued to track just above the upper prediction bound. In **Week 47** (20 26 Nov 2022), there were **578** excess deaths with a p-score of 15%. The cumulative total number of excess natural deaths in this age group since **3 May 2020** is about **251,200**.

Week	Date	Weekly excess deaths from natural causes for persons 60+	p-score
		years	
37	11-Sep-22 – 17-Sep-22	588	14.0%
38	18-Sep-22 – 24-Sep-22	591	14.3%
39	25-Sep-22 – 1-Oct-22	769	19.1%
40	2-Oct-22 – 8-Oct-22	796	18.9%
41	9-Oct-22 – 15-Oct-22	689	17.0%
42	16-Oct-22 – 22-Oct-22	478	12.2%
43	23-Oct-22 – 29-Oct-22	419	10.8%
44	30-Oct-22 – 5-Nov-22	669	16.8%
45	6-Nov-22 – 12-Nov-22	646	16.4%
46	13-Nov-22 – 19-Nov-22	714	18.6%
47	20-Nov-22 – 26-Nov-22	578	15.2%

- The uptick in natural deaths experienced in Week 40 (2 − 8 Oct 2022) in several provinces has subsided with the numbers in all provinces now falling close to the upper prediction bound or clearly within their prediction bounds.
- Per capita excess death rates have been calculated for the provinces to scale the cumulative deaths for the population size of each province (**Table 1**). By the end of **Week 47** (20 26 Nov 2022), the national excess death rate since 3 May 2020 was **567** per **100,000** population.
- The provinces with the highest cumulative numbers of excess deaths at the end of Week 47 (20 26 Nov 2022), are, in order, KwaZulu-Natal, Gauteng and Eastern Cape. The ranking changes to Eastern Cape, Northern Cape and Free State for the crude death rates per capita (i.e., taking size of the provincial populations into account) and to Northern Cape, Eastern Cape and KwaZulu-Natal using the agestandardised rates per capita (i.e., taking into account the age distribution of the provincial population). The Western Cape, followed by Gauteng, continue to have the lowest cumulative age standardised per capita rates.
- The number of unnatural deaths reached a high of 1,557 in Week 15 (10 16 Apr 2022), the week of the floods in KwaZulu-Natal, and reverted to the predicted monthly cyclical pattern thereafter. After tracking within the prediction bounds for several weeks, numbers increased to the upper prediction bound in Weeks 39 & 40, thereafter falling closer to the predicted number with 1,188 deaths in Week 47 (20 26 Nov 2022).



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations


Table 1: Number of excess natural deaths of persons by province and metro relative to revised predicted number based on the observed drop during lockdown, South Africa 2020/22


Region	Period	Excess deaths vs revised base	Excess deaths per 100,000 population	Age standardised excess death rate per 100,000
South Africa	3 May 20 – 26 Nov 22	337,747	567	567
Province				
Eastern Cape	31 May 20 – 26 Nov 22	58,533	889	717
Free State	21 Jun 20 – 26 Nov 22	19,496	670	670
Gauteng	7 Jun 20 – 26 Nov 22	66,383	425	468
KwaZulu-Natal	7 Jun 20 – 26 Nov 22	69,688	609	701
Limpopo	21 Jun 20 – 26 Nov 22	37,868	641	561
Mpumalanga	21 Jun 20 – 26 Nov 22	25,916	539	580
Northern Cape	28 Jun 20 – 26 Nov 22	10,318	881	825
North West	28 Jun 20 – 26 Nov 22	18,334	455	467
Western Cape	3 May 20 – 26 Nov 22	31,212	442	389
Metropolitan Municipality				
Buffalo City	31 May 20 – 26 Nov 22	7,655		
City of Cape Town	3 May 20 – 26 Nov 22	24,515		
Ekurhuleni	7 Jun 20 – 26 Nov 22	14,936		
eThekwini	14 Jun 20 – 26 Nov 22	14,108		
Johannesburg	7 Jun 20 – 26 Nov 22	23,256		
Mangaung	21 Jun 20 – 26 Nov 22	6,225		
Nelson Mandela Bay	31 May 20 – 26 Nov 22	9,085		
City of Tshwane	7 Jun 20 – 26 Nov 22	12,090		


Note: Period has been determined based on when an upturn in the number of natural deaths became apparent. Parts do not sum to the whole because office closures due to Covid-19 may have led to registration of deaths at other offices which may not be in the same area, and random fluctuation at the point at which the baseline is determined.

Table 2: Number of excess deaths from all causes of persons by province and metro relative to predicted number based on historical trend, South Africa 2020/22


	Excess deaths vs	Excess deaths per	
Region	forecast	100,000 population	
South Africa	343,053	576	
Province			
Eastern Cape	61,669	937	
Free State	19,346	664	
Gauteng	63,735	409	
KwaZulu-Natal	73,811	645	
Limpopo	38,538	652	
Mpumalanga	25,772	536	
Northern Cape	10,254	876	
North West	17,782	442	
Western Cape	32,145	455	
Metropolitan Municipality			
Buffalo City	8,342		
City of Cape Town	23,342		
Ekurhuleni	13,767		
eThekwini	13,238		
Johannesburg	23,834		
Mangaung	7,109		
Nelson Mandela Bay	9,230		
City of Tshwane	12,736		











Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations


















































Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations. As only a quarter to a third of unnatural deaths in the most recent week are processed at the time of the survey, the estimate for the most recent week is quite uncertain.

#### Vertical lines in order

- 0 Week Disaster Management Act implemented
- 1 Week lockdown level 5 introduced
- Week lockdown changed to level 4, with curfew
- 3 Week lockdown changed to level 3 including unbanning of alcohol
- 4 Week alcohol re-banned, and a curfew re-introduced
- 5 Week lockdown changed to level 2, including unbanning of alcohol
- 6 Week lockdown changed to level 1
- 7 Week lockdown changed to level 3 advanced (re-banning alcohol and a extension of curfew)
- 8 Week lockdown relaxed to allow sale of alcohol 4 days/week and reduce curfew
- 9 Week lockdown relaxed to allow sale of alcohol except during curfew and reduce curfew to midnight to 4am
- 10 Week lockdown changed to level 3 advanced (limiting alcohol and a extending of curfew)
- 11 Week lockdown changed to level 4, with re-banning of alcohol, curfew 9pm-4am
- 12 Week of unrest in KZN and GT
- 13 Week lockdown changed to level 3 advanced (alcohol 4 days/w, curfew 10pm-4am)
- 14 Week lockdown changed to level 2 advanced (alcohol 5 days/w, curfew 11pm-4am)
- 15 Week lockdown changed to level 1 advanced (no alcohol post 11pm, curfew 12pm-4am, large gatherings)
- 16 Week lockdown level 1 advanced (removed limits on alcohol & curfew, allowed larger gatherings)
- 17 Week lockdown level 1 advanced (allowed full school attendance, reduced isolation & quarantine requirements)
- 18 Week lockdown level 1 advanced (no masks outdoors, larger gathering permitted, no travel testing for vaccinated)
- 19 KZN floods
- 20 All COVID restrictions lifted on 22 June