

REPORT ON WEEKLY DEATHS IN SOUTH AFRICA

1 JANUARY – 21 JULY 2020
(WEEK 29)

Debbie Bradshaw, Ria Laubscher,
Rob Dorrington, Pam Groenewald, Tom Moultrie

Burden of Disease Research Unit
South African Medical Research Council
28 July 2020



UCT Centre
for Actuarial
Research

Warning: The Department of Home Affairs has faced sporadic temporary office closures, particularly in areas that are more affected by COVID-19. This may affect our allocation of a death to a metro area. For example, a death that occurred in the City of Cape Town might have been registered at an office outside of the City because of a temporary closure. Closure may also cause a delay in the processing of the death registration which would result in an underestimate of the deaths in the most recent week. This accounts for the kinks in what should otherwise be a smooth increase in numbers of deaths in Cape Town and Buffalo, for example.

Glossary:

Actual number of deaths: The actual number of deaths in South Africa have been estimated from the numbers recorded on the National Population Register using weighting factors set to produce results consistent with those of the annual Rapid Mortality Surveillance Report to account for deaths of persons who are not on the National Population Register as well as those that have not been registered with the Department of Home Affairs. The estimated number of weekly deaths can be downloaded in Excel with this report from the SAMRC website: <https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa>

Excess deaths: There is no universal definition of, or understanding of what is meant by, “excess mortality”. It is a term used in epidemiology and public health that refers to the number of deaths that are occurring above what we would normally expect. The WHO uses the term to describe “Mortality above what would be expected based on the non-crisis mortality rate in the population of interest. Excess mortality is thus mortality that is attributable to the crisis conditions. It can be expressed as a rate (the difference between observed and non-crisis mortality rates), or as a total number of excess deaths.”

Excess natural deaths associated with COVID-19: Generally, the number of excess deaths per week is calculated as the number of all-cause deaths in that week less the number that might be assumed to have occurred had there not been the epidemic (i.e. the counterfactual number), provided that the counterfactual is lower. However, this approach has generally only been applied to countries where deaths have been tracking the counterfactual before the onset of significant numbers of COVID-19 related deaths. The method provides a poor estimate of the numbers of COVID-19 and collateral deaths in the early stages of the epidemic when this is not the case. Thus, we estimate the numbers of COVID and collateral deaths, once a clear upward trend is evident, as the number of actual deaths less a baseline number determined as a proportion of the predicted number. The proportion is calculated such that the excess deaths in that week is equal to the confirmed number of COVID-19 deaths for that week. The cumulative number of excess deaths comprises the sum of the weekly excess plus the cumulative number of confirmed COVID deaths prior to the establishment of the clear upward trend. Where there is no clear indication of an upward trend, we have not calculated excess deaths. It is important to note that this estimate of the number of excess deaths is an estimate of the number of deaths in excess of expectation, due to the Covid-19 epidemic and not of those infected with the SARS-CoV-2 virus alone (i.e. it includes incidental deaths resulting from such things as shortage of health care and medications due to either the demands on the health systems by the virus or strategies to combat the epidemic).

Data Source


Basic demographic information for all deaths registered on the National Population Register are provided to the SAMRC on a weekly basis. Since the number of deaths has a seasonal trend, historical data from 2018 and 2019 have been used to predict the number of deaths that could be expected during 2020. Before this was done, the deaths were weighted to account for incomplete registration of deaths and those that do not have a South African ID number. The weights were calculated by age, sex, metro/non-metro and natural/unnatural cause to be consistent with the weights applied in the annual Rapid Mortality Surveillance Reports.¹

While we have built up a good sense of the adjustment at a national level through the annual RMS reports and the National Burden of Disease Study, estimating completeness of registration of deaths below national level is challenging, particularly given limitations of data available to inform such an exercise, and has required numerous assumptions. Thus, the resulting estimates need to be treated with a degree of caution.

The Excel forecast function² has been used to predict values for each week of 2020 based on a linear annual trend, allowing for a seasonal effect over the year. In addition, 95% prediction intervals have been estimated for the predicted weekly number of deaths for 2020 to give a basis to assess fluctuations. The forecasts have been applied to the estimated actual number of deaths.

Graphs of the estimated weekly number of deaths, classified by date of occurrence, not reporting, up until epidemiological **week 29** (i.e. the period from **1 January 2020** till **21 July 2020**) based on the data received on 27 July 2020 are shown below. *The figures plot the estimated numbers of deaths at the start date of each week.* Data for the most recent week has been scaled up to account for the lag in processing registrations. Based on previous data, the numbers at the national level have been increased by 5.5%.

Numbers of weekly deaths for persons 1+ years from all causes from the National Population Register are shown in the figure below. It can be seen that in June, the average number would be as forecast. However, there is a clear increase from being lower at the beginning of the month to being somewhat higher by the end of the month. The trend continues into July.



¹ Dorrington RE, Bradshaw D, Laubscher R, Nannan N (2020). Rapid mortality surveillance report 2018. Cape Town: South African Medical Research Council.

² The Excel function implements is the Holt-Winters triple exponential smoothing (the AAA sub-method).

Sub-national statistics have been compiled for the provinces and metros by allocating the deaths according to the Home Affairs office where the death was registered. It is assumed that most of the deaths within an area are registered at an office in the same area. The numbers of deaths from **natural causes** are reported for each province and each of the metros.

Estimating the number of COVID-19 and collateral (i.e. deaths arising as a result of the impact of SARS-COV-2 epidemic and the management of this on the provision of health care) deaths is not straight-forward. In developed countries this is measured using the extent which deaths due to all causes exceed what might be expected had the epidemic not occurred. For various reasons this is not the best way of estimating COVID and collateral deaths in South Africa, particularly in the early weeks of the epidemic.

For a start, unnatural deaths are a higher proportion of all deaths in South African than in developed countries, and reduced significantly as economic and social activity became more limited, particularly in lockdown stage 5. Thus, it was decided to limit our focus to the numbers of deaths from natural causes.

Further, it was observed that during lockdown the number of deaths due to natural causes fell well below the predicted number. The weekly numbers tracked the lower prediction interval, probably due to the impact of lockdown and non-pharmaceutical interventions (NPIs) on the spread of non-COVID communicable diseases.

Thus, using the predicted value as the base would understate the impact of the COVID-19 epidemic, identify no COVID or collateral deaths until the week starting 17 June, when there were already over 2,000 confirmed COVID deaths. It was therefore decided to take as the expected number of deaths a proportion of the predicted number, that matched the proportion³ in the week prior to week when there was a clear rapid increase in numbers (e.g. the week starting on 6 May for Cape Town, Western Cape and nationally), and the number of cumulative COVID and collateral cases was set to the number of confirmed cases for that week.

Continuing to use this adjusted base until late June is supported by the observation that the number of natural deaths in the provinces less affected by COVID-19 continued to track lower than predicted numbers through to the end of June. However, following the emergence of COVID-19 in most of the provinces, and with uncertainty about the counterfactual following the seasonal winter peak in numbers of deaths, the baseline is transitioning to the predicted number of deaths by next week. The estimated numbers of excess natural deaths are reported in **Table 1**, relative to this revised base. These numbers are indicated on the graphs. It is important to point out that although it is probable that the bulk of these numbers of 'excess deaths' are likely to be due to COVID-19 it is unclear at this stage what proportion are COVID deaths and what proportion are collateral deaths.


Although less adequate as a measure at the early stage of the epidemic, **Table 2**, provides estimates of 'excess mortality' using the measure employed more generally based on all-cause deaths.

Births were not registered by the Department of Home Affairs during lockdown stage 5. This means any that die before the backlog is processed will not be placed on the National Population Register and thus that the deaths of these births will not be captured. **This report presents the estimated weekly deaths of persons 1 year and older.** Registered births are again being added to the Population Register, but either there remains a backlog in processing or a lower proportion of births are being registered since lockdown. Once we have confidence that registration of deaths is back to previous levels we will include deaths under age 1.

³ With a small adjustment to ensure that the number of COVID deaths matched the numbers of confirmed cases for that week.

Trends

- The all-cause national number of deaths of persons 1+ years of age per week has risen to **16,223** and is **63%** higher than the predicted number based on historical data in the week ending **21 July 2020**. When compared with the predicted numbers, there was an excess of **6,256** deaths in the latest week.
- The number of deaths from natural causes is also significantly higher than the predicted number, for persons 1-59 years and 60+ years.
- In the period, **6 May - 21 July 2020**, there has been an excess of **22,279** deaths from natural causes of persons 1+ year old when using the revised base accounting for lower mortality during lockdown. For people 1-59 years the excess is **7,536** and **14,697** for people 60+ years.
- **Table 1** shows the estimated excess number of natural deaths in metro areas and the provinces. Deaths from natural causes in the **City of Cape Town (3,271)**, **Johannesburg (2,462)**, **Ekurhuleni (2,033)**, **Nelson Mandela Bay (1,316)**, **Buffalo City (847)** and **City of Tshwane (673)** metros continued increasing in the week up to **21 July 2020**.
- **Free State, Mpumalanga, Northern Cape and North West** have joined **Western Cape, Eastern Cape, Gauteng** and **KwaZulu-Natal** and are also experiencing significantly more natural deaths than predicted. Compared with the predicted number of natural deaths from historical data in the week ending **21 July 2020**, **Gauteng** had **127%** more, **Eastern Cape** had **118%** more, Free State had **92%** more, **KwaZulu-Natal and Northern Cape** had **69%** more, **North West** had **58%** more, **Mpumalanga** had **42%** more and **Western Cape** had **34%** more.
- The number of deaths from unnatural causes (e.g. road traffic fatalities and homicides) was **29%** below the predicted number for the week ending **21 July 2020**.



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations




Table 1: Number of excess natural deaths of persons 1+ years by province and metro relative to revised predicted number based on the observed drop during lockdown, South Africa 2020

Region	Period	Excess deaths vs revised base
South Africa	6 May – 21 July	22,279
Province		
Eastern Cape	3 June – 21 July	6,411
Free State	24 June – 21 July	752
Gauteng	10 June – 21 July	6,620
KwaZulu-Natal	6 May – 21 July	2,632
Limpopo	24 June – 21 July	527
Mpumalanga	24 June – 21 July	627
Northern Cape	14 July – 21 July	164
North West	1 July – 21 July	566
Western Cape	6 May – 21 July	4,133
Metropolitan Municipality		
Buffalo City	3 June – 21 July	847
City of Cape Town	6 May – 21 July	3,271
Ekhuruleni	10 June – 21 July	2,033
Ethikweni	17 June – 21 July	673
Johannesburg	10 June – 21 July	2,462
Mangaung		-
Nelson Mandela Bay	3 June – 21 July	1,316
City of Tshwane	10 June – 21 July	865




Note: Period has been determined based on when an upturn in the number of natural deaths became apparent. Parts do not sum to the whole because office closures due to Covid-19 may have led to registration of deaths at other offices which may not be in the same area, and random fluctuation at the point at which the baseline is determined.

Table 2: Number of excess deaths from all causes of persons 1+ years by province and metro relative to predicted number based on historical trend, South Africa 2020




Region	Excess deaths vs forecast
South Africa	16,954
Province	
Eastern Cape	5,932
Free State	731
Gauteng	6,102
KwaZulu-Natal	2,046
Limpopo	205
Mpumalanga	584
Northern Cape	210
North West	642
Western Cape	2,983
Metropolitan Municipality	
Buffalo City	645
City of Cape Town	2,288
Ekhuruleni	1,709
Ethikweni	583
Johannesburg	2,334
Mangaung	82
Nelson Mandela Bay	1,279
City of Tshwane	852






Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations






Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations






Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations





Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations




Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations



Vertical time lines:

1. Week lockdown level 5 introduced
2. Week lockdown changed to level 4 with curfew
3. Week lockdown changed to level 3 including unbanning sale of alcohol
4. Week sale of alcohol re-banned and curfew re-introduced

Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations